

DATE: 2/21/2017

FARM: RUNNYMEDE FARM

NAME: AVARICITY 16

DOCTOR: DH Rodgerson, DVM, MS, Diplomate ACVS

LF Fetlock: NSF*

RF Fetlock: NSF*

LH Fetlock: NSF*

RH Fetlock: NSF*

Left Knee: NSF*

Right Knee: NSF*

Left Hock: Small spur or irregular margin to the front of lower hock joints. May not change much by sale time. Mild tarsitis.

Right Hock: NSF*

Left Stifle: NSF*

Right Stifle: Lucency on medial condyle of femur. Reshoot in 8 weeks. You can continue on chondroprotective agents (Reflex and/or Adequan). Doers look better and appears to be more dense then previous. There is a shallow defect in subchondral bone. If sound I would not inject again.

** NSF= No significant radiographic findings.*

Comments: Mild tarsitis is a small amount of bone proliferation, spur, or mild changes to lower hock joints. Typically not discount too much if changes are mild. Cyst is the main issue but if horse stays sound, the long term prognosis maybe okay.

The above expresses the opinion of the examiner based on results and findings by the examiner at the time of the radiographic examination. This opinion is rendered only to the party requesting said examination and may not be relied upon by any other party. Results of examinations of this nature and the opinion expressed shall not carry with it any warranty of guarantee of any other kind, expressed or implied.